



### Zolpidem - General Effects

**Zolpidem** (Ambien) is a sedative-hypnotic and central nervous system (CNS) depressant.

1. Zolpidem is used as a hypnotic agent and for the short-term treatment of insomnia.
  - a. Zolpidem has a half-life of 1.4-4.5 hours.
  - b. Zolpidem has a general therapeutic range, but experienced effects can vary based on an individual’s prescription history.
    - i. Therapeutic range refers to the blood concentration expected to achieve the desired therapeutic effects. Due to many factors such as prescription history, dosage, tolerance, drug-drug interactions and use, an individual may exhibit signs of impairment even though blood concentrations fall within the therapeutic range.
2. General effects of Zolpidem include but are not limited to: drowsiness, dizziness, amnesia, headache, and nausea.
  - a. Because zolpidem acts as a sleep inducer, therapeutic and above-therapeutic doses of zolpidem can have a negative impact on driving. Some of the possible effects of zolpidem use include slowed reaction time, erratic lane traveling, disorientation, slurred speech, lack of balance, and falling asleep at the wheel.
3. The longer an individual uses a drug, the more they can build up a tolerance to its effects. Tolerance occurs when an individual no longer responds to the drug in the way that they initially responded. When an individual gains tolerance to a drug, a higher dose of the drug is necessary to achieve the same level or response initially achieved. As tolerance is gained, it may reduce some of the possible negative effects of a drug.
4. Drug metabolism (alcohol excluded) exhibits first order kinetics, or the elimination of a constant fraction of drug quantity per unit of time, which means that the amount eliminated is proportional to the drug concentration.
5. The use of more than one drug at a time may enhance the effects the drugs would otherwise have on their own, leading to greater impairment.

**References**

1. Baselt, R.C. (2014). Disposition of toxic drugs and chemicals in man. Biomedical Publications, Seal Beach, CA.
2. Moffat, A. C., Osselton, M. D., Widdop, B., & Watts, J. (2011). Clarke's analysis of drugs and poisons (3rd ed., Vol. 1). Pharmaceutical Press.
3. Levine, B., & Vina Spiehler. (2020). Pharmacokinetics and Pharmacodynamics. In B. Levine (Ed.), Principles of Forensic Toxicology (4th ed., pp. 77–93). essay, AACC Press.
4. Logan, B.K., Couper, F.J. (2000) Zolpidem and Driving Impairment. Journal of Forensic Sciences. 46:1, 105-110.

Template and general factual statements used from Colorado Bureau of Investigation’s monograph system. Edits and fact checking performed by Wisconsin State Crime Laboratory prior to publication.

\*\* The interpretive information provided is not exhaustive nor meant to encompass all scenarios where toxicological results are reported. Interpretive information is meant to serve as a general guide for the reader and that for any given case, consultation with a forensic toxicologist is recommended. \*\*

| — Zolpidem Monograph |          |                       |           |             |
|----------------------|----------|-----------------------|-----------|-------------|
| Document ID          | Revision | Effective Date        | Status    | Page Number |
| 50146                | 1        | 12/20/2024 1:01:36 PM | Published | 1 of 1      |